



Every effort has been made to ensure the accuracy of the information published. However, it remains the responsibility of the readers to familiarize themselves with the product information contained on the US product label or package insert.

---

# SCALIBOR<sup>®</sup> PROTECTOR BAND FOR DOGS

*Intervet/Schering-Plough Animal Health*

*Flea & Tick Collar for Dogs*

**LONG LASTING PROTECTION**

*For up to 6 months*

**PATENTED**

*Insecticide-Release Technology*

**KILLS FLEAS**

**KILLS TICKS**

*Including deer ticks which may carry Lyme Disease*

**WATER RESISTANT**

**ADJUSTABLE**

*One Size Fits All*

**READ ENTIRE LABEL BEFORE EACH USE**

**USE ONLY ON DOGS**

Deltamethrin 4% Collar provides full season, up to 6 months protection against fleas and ticks

**PRECAUTIONARY STATEMENTS**

**Hazards to Humans and Domestic Animals**

**CAUTION:**

Do not open protective pouch until ready to use. Do not let children play with this collar.

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling.

Do not use on puppies under 12 weeks. Consult a veterinarian before using this product on debilitated, aged, pregnant, medicated or nursing animals. Sensitivities may occur after using ANY pesticide products for pets. If signs of sensitivity occur, remove collar and bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

This collar is intended for use only as an insecticide generator and is not to be taken internally by man or animals. Applying other pesticides on the dog may not be necessary while the collar is being worn.

**DIRECTIONS FOR USE:**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Deltamethrin 4% Collar, containing deltamethrin insecticide, has been specially formulated using patented insecticide-release technology. Maximum effectiveness may not occur for 2-3 weeks after collar placement. Fleas (*Ctenocephalides* sp.) on the dog will be killed and ones which are present in the dog's environment that may appear on your pet will be killed. Collar will also kill ticks for up to 6 months. Collar will kill ticks including Brown dog tick (*Rhipicephalus sanguineus*), American dog tick (*Dermacentor variabilis*), and deer ticks (*Ixodes scapularis* and *Ixodes pacificus*) which may carry the Lyme disease. This collar should be worn continuously. Reapply a new collar every 6 months.

Place the collar around dog's neck, buckle and adjust for proper fit. Cut off approximately 2 inches from the buckle and dispose of excess length by wrapping in newspaper and placing in trash. The collar must be worn loosely so that two fingers may be placed between collar and dog's neck. Living and rest areas of pet must also be treated with appropriate pest control measures to ensure control of pests. Wetting will not impair the collar's effectiveness of the pet's protection. If the dog goes swimming or is out in the rain, it is not necessary to remove the collar.

Deltamethrin 4% Collar may be used in addition to a lead or constraint collar. Use only one Deltamethrin 4% Collar

at a time.

**FIRST AID**

**IF SWALLOWED**

- Call a poison control center or doctor immediately for treatment advice
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by the poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

**IF ON SKIN OR CLOTHING**

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

**IF IN EYES**

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

For incidents involving **HUMANS**, call 1-800-680-9206

For incidents involving **ANIMALS**, call 1-800-548-2423

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

**STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage and disposal.

**Pesticide Storage:** Store this product in a cool, dry place away from children and pets.

**Pesticide and Container Disposal:** Do not reuse container or used collar. Dispose of in trash.

ACTIVE INGREDIENT:	Percentage by Weight
Deltamethrin	4.0%
INERT INGREDIENTS	96.0%
Total	100.0%

EPA Reg No. 68451-1

EPA Est. No. 68451-FRA-1

NET CONTENTS: 1 Collar

NET WT. 0.9 oz

Distributed by: Intervet Inc., Roseland, NJ 07068

Product of France

**NAC No.:** 10473010