

**DURVET, INC.****100 S.E. MAGELLAN DRIVE, BLUE SPRINGS, MO, 64014**

Telephone: 816-229-9101

Toll-Free: 800-821-5570

Fax: 816-224-3080

Website: [www.durvet.com](http://www.durvet.com)Email: [info@durvet.com](mailto:info@durvet.com)

Every effort has been made to ensure the accuracy of the information published. However, it remains the responsibility themselves with the product information contained on the USA product label or package insert.

**CANINE SPECTRA®5*****Durvet*****Canine Distemper-Adenovirus Type 2-Parainfluenza-Parvovirus Vaccine**


Modified Live Virus

**FOR USE IN DOGS AND PUPPIES****KEEP REFRIGERATED**

**Description:** Spectra®5 vaccine is a combination of antigenic, attenuated strains of Canine Distemper, Canine Adenovirus Type 2 (CAV-2), Canine Parainfluenza, and Canine Parvovirus Type 2b, propagated in cell line tissue cultures. The CAV-2 fraction cross-protects against respiratory infection caused by infectious canine hepatitis (CAV-1).

**Indications:** For the vaccination of healthy, susceptible dogs and puppies as an aid in the reduction of diseases caused by canine distemper, canine adenovirus types 1 and 2, parainfluenza, and parvovirus.

**Directions and Dosage:** The dosage is 1 mL injected intramuscularly or subcutaneously. Open syringe by twisting or tapping the cap against a hard surface to break the heat weld. Prepare the vaccine by injecting the diluent into the vial containing the desiccated vaccine cake. Shake well. Withdraw entire contents into the syringe. Thoroughly prepare injection site with antiseptic. Subcutaneous administration: Lift the loose skin behind the neck or behind front leg and insert needle. Inject entire contents of syringe. Do not inject directly into blood vessel (see note below). Intramuscular administration: Insert needle into muscle of the hind limb (see illustration). Inject entire contents of syringe. Do not inject directly into blood vessel. Important note: Before injecting vaccine pull back slightly on syringe plunger. If blood enters the syringe freely, choose another injection site. Puppies 9 weeks or younger: Vaccinate healthy puppies at 3-4 week intervals until 16 weeks of age for at least 3 doses. Puppies over 9 weeks and adults: Give a minimum of 2 doses at 3-4 week intervals. Annual revaccination with a single dose is recommended.



**Storage:** Store out of direct sunlight at 35°-45°F (2°-7°C). Do not freeze. **Disposal:** Burn containers and unused contents.

**Precautions:** Do not vaccinate pregnant animals. Do not use in ferrets or mink. Protective immunity may not be completely established in all puppies vaccinated at less than 16 weeks of age because of maternal antibody interference. Only vaccinate healthy animals. Animals incubating any disease or stressed due to shipping, malnutrition, or parasitism may not achieve or maintain an adequate immune response. Although rare, severe allergic reactions (anaphylactoid) may occur that require immediate veterinary care.

**Preservatives:** Gentamicin and Amphotericin B.

**Antidote:** Epinephrine.

MADE IN THE USA

Produced by: Boehringer Ingelheim Vetmedica, Inc., St. Joseph, MO 64506 U.S.A.

U.S. Vet. License No. 124

Distributed by: Peak Marketing, P.O. Box 279, Blue Springs, MO 64013 U.S.A.

Spectra® 5 is a trademark of Peak Marketing LLC

One Dose with Syringe	Package contains a one dose vial of vaccine, a one mL vial of vaccine diluent, & one disposable syringe
25 Doses	25 (1 dose) vials of dry vaccine and 25 (1 mL) vials of vaccine diluent

**CPN:** 1084285.2