

**durvet****Duramycin 72-200****(Oxytetracycline)  
200 mg/mL  
Antibiotic**

Each mL contains 200 mg of oxytetracycline base  
as amphoteric oxytetracycline.  
For the treatment of disease in beef cattle; dairy cattle;  
calves, including preruminating (veal) calves; and swine.

ANADA 200-306, Approved by FDA

**Net Contents: 250 mL**

Duramycin 72-200 (oxytetracycline) is a sterile, preconstituted solution of the broad-spectrum antibiotic oxytetracycline. Each mL contains 200 mg of oxytetracycline base as oxytetracycline amphoteric, and on a w/v basis, 40.0% 2-pyrrolidone, 5.0% povidone, 1.8% magnesium oxide, 0.2% sodium formaldehyde sulfoxylate (as a preservative), monoethanolamine and/or hydrochloric acid as required to adjust pH.

**CAUTION:** When administered to cattle, muscle discoloration may necessitate trimming of the injection site(s) and surrounding tissues during the dressing procedure.

**WARNINGS:** Discontinue treatment at least 28 days prior to slaughter of cattle and swine. Milk taken from animals during treatment and for 96 hours after the last treatment must not be used for food.

**PRECAUTIONS:** Exceeding the highest recommended level of drug per lb of body weight per day, administering more than the recommended number of treatments, and/or exceeding 10 mL intramuscularly or subcutaneously per injection site in adult beef and dairy cattle, and 5 mL intramuscularly per injection site in adult swine, may result in antibiotic residues beyond the withdrawal period.

TAKE TIME

OBSERVE LABEL  
DIRECTIONS

Lot No.:

Exp. Date:

Rev. 09-11  
023215L03**DOSAGE:**

**CATTLE:** A single dosage of 9 mg of oxytetracycline per lb of body weight (4.5 mL/100 lb) administered intramuscularly or subcutaneously is recommended in the treatment of the following conditions:

- (1) bacterial pneumonia caused by *Pasteurella* spp. (shipping fever) in calves and yearlings, where retreatment is impractical due to husbandry conditions, such as cattle on range, or where repeated restraint is inadvisable.
- (2) infectious bovine keratoconjunctivitis (pink eye) caused by *Moraxella bovis*.

**SWINE:** A single dose of 9 mg of oxytetracycline per lb of body weight (4.5 mL/100 lb) administered *intramuscularly* is recommended in the treatment of bacterial pneumonia caused by *Pasteurella multocida* in swine, where retreatment is impractical due to husbandry conditions or where repeated restraint is inadvisable.

Refer to Package Insert for Complete Directions.

**STORAGE:** Store at 59° to 86°F (15° to 30°C).  
Keep from freezing.

**Not for Human Use. For Animal Use Only.**

**Restricted Drug - California. Use Only as Directed.**

**Manufactured for:**

**DURVET, INC.**

Blue Springs, Missouri 64014

www.durvet.com

**Made in the UK.**

**Manufactured by:**

Norbrook Laboratories Limited,

Newry, BT35 6PU, Co. Down, Northern Ireland.



6

7

Net Contents: 250 mL  
 ANADA 200-306, Approved by FDA  
 For the treatment of disease in beef cattle; dairy cattle; calves, including preruminating (veal) calves; and swine.  
 Each mL contains 200 mg of oxytetracycline base as amphoteric oxytetracycline.



**durvet**

NDC 30798-237-13

NDC 30798-237-13

**durvet**

**Duramycin 72-200**



**(Oxytetracycline)  
 200 mg/mL  
 Antibiotic**

Each mL contains 200 mg of oxytetracycline base as amphoteric oxytetracycline.

**For the treatment of disease in beef cattle; dairy cattle; calves, including preruminating (veal) calves; and swine.**

ANADA 200-306, Approved by FDA

**Net Contents: 250 mL**

Duramycin 72-200 is a sterile, preconstituted solution of the broad-spectrum antibiotic oxytetracycline.

**CAUTION:**  
 When administered to cattle, muscle discoloration may necessitate trimming of the injection site(s) and surrounding tissues during the dressing procedure.

**WARNINGS:**  
 Discontinue treatment at least 28 days prior to slaughter of cattle and swine. Milk taken from animals during treatment and for 96 hours after the last treatment must not be used for food. Rapid intravenous administration may result in animal collapse. Oxytetracycline should be administered intravenously slowly over a period of at least 5 minutes.

**PRECAUTIONS:**  
 Exceeding the highest recommended level of drug per lb of body weight per day, administering more than the recommended number of treatments, and/or exceeding 10 mL intramuscularly or subcutaneously per injection site in adult beef cattle and dairy cattle, and 5 mL intramuscularly per injection site in adult swine, may result in antibiotic residues beyond the withdrawal period.

Refer to Package Insert for Complete Directions

**Storage:**  
 Store at 59° to 86°F (15° to 30°C). Keep from freezing.

Restricted Drug - California. Use Only as Directed.

Not for Human Use. For animal use only.

**Manufactured for:**  
**DURVET, INC.**  
 Blue Springs, Missouri 64014  
 www.durvet.com

Made in the U.K.

**Manufactured by:**  
 Norbrook Laboratories Limited  
 Newry, BT35 6PU, Co. Down,  
 Northern Ireland

TAKE TIME



**OBSERVE LABEL DIRECTIONS**

NDC 30798-237-13

**durvet**

**Duramycin 72-200**



**(Oxytetracycline)  
 200 mg/mL  
 Antibiotic**

Each mL contains 200 mg of oxytetracycline base as amphoteric oxytetracycline.

**For the treatment of disease in beef cattle; dairy cattle; calves, including preruminating (veal) calves; and swine.**

ANADA 200-306, Approved by FDA

**Net Contents: 250 mL**

**CATTLE DOSAGE GUIDE**

At the first signs of pneumonia or pinkeye, \* administer a single dose of Duramycin 72-200 (200 mg/mL) by deep intramuscular injection, or subcutaneously, according to the following weight categories. \*\*

Animal Weight (lb)	Number of mL or cc
100	4.5
200	9.0
300	13.5
400	18.0
500	22.5
600	27.0
700	31.5
800	36.0
900	40.5
1000	45.0
1100	49.5
1200	54.0

\* See package insert for dosing instructions for other indicated diseases and full product information.  
 \*\* Do not administer more than 10 mL at any one injection site (1-2 mL per site in small calves).  
 Discontinue treatment at least 28 days prior to slaughter.

**SWINE DOSAGE GUIDE**

At the first signs of pneumonia, \* administer Duramycin 72-200 (200 mg/mL) by deep intramuscular injection, according to the following weight categories. \*\*

Animal Weight (lb)	Number of mL or cc
10	0.5
25	1.1
50	2.3
75	3.4
100	4.5
125	5.6
150	6.8
175	7.9
200	9.0
225	10.1
250	11.3
275	12.4
300	13.5
325	14.6

\* See package insert for dosing instructions for other indicated diseases and full product information.  
 \*\* Do not administer more than 5 mL at any one injection site.  
 Discontinue treatment at least 28 days prior to slaughter.

Lot No.:  
 Exp. Date:

Rev. 09-11  
 6215302C02

