



Compendium of Veterinary Products - US edition

WG CRITICAL CARE, LLC
120 ROUTE 17 NORTH, PARAMUS, NJ, 07652
Customer Service Tel.: 888-493-0861
Customer Service Fax: 414-434-6680
Customer Service Email: CustomerService@wgccrx.com
Website: www.wgcriticalcare.com
Email: info@wgccah.com



Every effort has been made to ensure the accuracy of the information published. However, it remains the responsibility of the readers to familiarize themselves with the product information contained on the USA product label or package insert.

MASTI-CLEAR[®]

WG Critical Care

(penicillin G procaine in sesame oil)

100,000 units/10mL Syringe

For Intramammary Infusion in Lactating Cows Only

Antibacterial

NDC 10515-014-10

DESCRIPTION:

Each syringe contains:

100,000 units of penicillin G procaine, sesame oil 8.7g, calcium carbonate 0.089g, corn starch 0.089g, hydrogenated castor oil 0.044g.

INDICATIONS

This product is intended for the treatment of bovine mastitis in lactating cows. This product is effective against udder infection caused by the following susceptible microorganisms:

Streptococcus agalactiae, *Streptococcus dysgalactiae* **and** *Streptococcus uberis*.

For the best results, it should be used promptly at the first signs of mastitis.

Signs of mastitis include:

1. **Inflammation of the udder or teat.** Inflammation may be recognized by redness, swelling, elevated temperature, or pain in the affected part.
2. **Abnormal milk.** Presence of flakes or clots in the milk, or milk of a watery character is usually indicative of a mastitis infection.
3. **High white-cell counts.** The routine use of the California Mastitis Test is encouraged. A high incidence of scores in excess of CMT 2 or a high CMT score on bulk milk samples may indicate a potential mastitis problem. In these situations a veterinarian should be consulted for definitive diagnosis and recommendations.

DIRECTIONS FOR USE

Dosage: The dose is 10mL of this product per infected quarter administered by intramammary infusion. Treatment may be repeated at intervals of 12 hours. Do not administer more than three consecutive doses.

Milk out udder completely. Wash udder and teats thoroughly with warm water containing a suitable dairy disinfectant. Dry thoroughly. The hands of the operator should be washed and dried before the administration of each treatment. Using the alcohol pads that are provided with each syringe wipe off teat, using a separate pad for each teat. Appropriate sanitation and management procedures to prevent and/or control bovine mastitis should be instituted.

When using this product remove plastic cover from tip of syringe and, while holding teat firmly, insert tip into streak canal; then push plunger and inject entire contents. After injection, grasp the end of the teat firmly, then gently massage the medication up the teat canal into the milk cistern.

Treated quarters should not be milked for at least six hours after treatment but should be milked at regular intervals thereafter. Treatment may be repeated at 12 hour intervals, up to a total of three doses, as indicated by the clinical response.

NOTE: Mastitis cannot be considered cured until bacteriologic examination of milk samples taken approximately three weeks after treatment demonstrates the absence of causative microorganisms.

WARNING:

1. Discard all milk for 60 hours (5 milkings) after the latest treatment.
2. Animals intended for human consumption must not be slaughtered within 3 days of the last treatment with Masti-Clear®.

3. Use of this product in a manner other than as directed in the labeling might result in violative residues.

PRECAUTIONS

Keep out of reach of children. Discard empty container; do not reuse. It is a violation of Federal law to use Masti-Clear® in food-producing animals other than as directed in the labeling. If abnormal milk, redness, or swelling persists or increases, discontinue use and consult your veterinarian.

USER SAFETY WARNINGS

Penicillins can cause allergic reactions in sensitized individuals. Topical exposure to penicillin may elicit mild to severe allergic reactions in some individuals. Repeated or prolonged exposure may lead to sensitization. Avoid direct contact of the product with the skin, eyes, mouth and clothing.

Persons with known hypersensitivity to penicillin should avoid exposure to this product.

In case of accidental eye exposure, flush with water for 15 minutes. In case of accidental skin exposure, wash with soap and water. Remove contaminated clothing. If allergic reaction occurs (e.g., skin rash, hives, difficult breathing), seek medical attention.

The material safety data sheet contains more detailed occupational safety information.

To report side effects in users, to obtain more information or to obtain a Material Safety Data Sheet, call 1-800-234-4263.

STORE AT CONTROLLED ROOM TEMPERATURE: 20° - 25°C (68° -77°F).

RESTRICTED DRUG (CALIFORNIA) - USE ONLY AS DIRECTED

THIS PRODUCT WAS MANUFACTURED BY A NON-STERILIZING PROCESS.

Manufactured By: G.C. Hanford Mfg. Co., Syracuse, NY 13201

NADA 65-081 APPROVED BY FDA

INS012LC 10 08/2011

NET CONTENTS:

10mL LBL012LC 09 04/11

CPN: 1865003.0

By using this site, you agree to the [Terms of Service](#) and [Privacy Policy](#).

[Contact Us](#) - [Logout](#)

Updated April 15, 2018

Copyright 2018 - Animalytix LLC