

Phoenix
 KC #350409
 Printed Side Up

INDICATIONS: Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism. IverCare® (ivermectin) Paste 1.87% provides effective treatment and control of the following parasites in horses.

Large Strongyles (adults) (also tissue stages), *S. equinus*, *T.odontophorus* spp., including *T. brevicauda* and *T. serratus*, and *Crateostomum acuticaudatum*, *Small Strongyles* (adults), including those resistant to some benzimidazole class compounds) – *Coronocylus* spp., including *C. coronatus*, *C. labiatus* and *C. labratus*, *Cyathostomum* spp., including *C. catinatum* and *C. pateratum*, *Cylicocyclus* spp., including *C. insignis*, *C. lepostomum*, *C. nassatus* and *C. brevicapsulatus*, *Cylicodonophorus* spp.; *Cylicostephanus* spp., including *C. callatus*, *C. goldi*, *C. longibivatus* and *C. minus*, and *Petrovinema pocalutum*, **Small Strongyles** (adults) – Fourth-stage larvae; **Pinworms** (adults and fourth-stage larvae) – *Oxyuris equi*; **Ascarids** (adults and third- and fourth-stage larvae) – *Parascaris equorum*; **Hairstworms** (adults) – *Trichostrongylus axei*; **Large-mouth Stomach Worms** (adults) – *Habronema muscae*; **Bots** (oral and gastric stages) – *Gasterophilus* spp., including *G. intestinalis* and *G. nasalis*; **Lungworms** (adults and fourth-stage larvae) – *Dicyocaulus amfeidi*, **Intestinal Threadworms** (adults) – *Strongyloides westeri*, *Strongyloides* caused by *Habronema* and *Draschia* spp. cutaneous third-stage larvae; Dermatitis caused by neck threadworm microfilariae, *Onchocerca* sp.

DOSEAGE AND ADMINISTRATION: This syringe contains sufficient paste to treat one 1500 lb horse at the recommended dose rate of 91 mcg ivermectin per lb (200 mcg/kg) body weight. Each weight marking on the syringe plunger delivers enough paste to treat 250 lb body weight. (1) While holding plunger, turn the knurled ring on the plunger 1/4 turn to the left and slide it so the side nearest the barrel is at the prescribed weight marking. (2) Lock the ring in place by making a 1/4 turn to cover from the tip of the syringe. (5) Insert the syringe tip into the horse's mouth at the space between the teeth. (6) Depress the plunger as far as it will go, depositing paste on the tip of the tongue. (7) Immediately raise the horse's head for a few seconds after dosing.

PARASITIC CONTROL PROGRAM: All horses should be included in a regular parasite control program with particular attention being paid to mares, foals and yearlings. Foals should be treated initially at 6 to 8 weeks of age, and routine treatment repeated as appropriate. Consult your veterinarian for a control program to meet your specific needs. IverCare® (ivermectin) Paste 1.87% effectively controls gastrointestinal nematodes and bots of horses. Regular treatment will reduce the chances of verminous arteritis caused by *Strongylus vulgaris*.

PRODUCT ADVANTAGES: Broad-spectrum Control. IverCare® (ivermectin) Paste 1.87% kills important internal parasites, including bots and the arterial stages of *S. vulgaris*, with a single dose. IverCare® (ivermectin) Paste 1.87% is a potent antiparasitic agent that is neither a benzimidazole nor an organophosphate.

ANIMAL SAFETY: IverCare® (ivermectin) Paste 1.87% may be used in horses of all ages, including mares at any stage of pregnancy. Stallions may be treated without adversely affecting their fertility.

WARNING: Do not use in horses intended for human consumption.

Not for use in humans. Keep this and all drugs out of reach of children.

Refrain from smoking and eating when handling. Wash hands after use. Avoid contact with eyes. The Material Safety Sheet (MSDS) contains more detailed occupational safety information. To report adverse reactions in users, to obtain more information, or to obtain a MSDS, contact Farnam Companies, Inc. at 800-234-2269.

PRECAUTIONS: IverCare® (ivermectin) Paste 1.87% has been formulated specifically for use in horses only. This product should not be used in other animal species as severe adverse reactions, including fatalities in dogs, may result.

Environmental Safety: Ivermectin and excreted ivermectin residues may adversely affect aquatic organisms. Do not contaminate ground or surface water. Dispose of the syringe in an approved landfill or by incineration.

INFORMATION FOR HORSE OWNERS: Swelling and itching reactions after treatment with Ivermectin Paste have occurred in horses carrying heavy infections of neck threadworm microfilariae (*Onchocerca sp.*). These reactions were most likely the result of microfilariae dying in large numbers. Symptomatic treatment may be advisable. Consult your veterinarian should any such reactions occur. Healing of summer sores involving extensive tissue changes may require other appropriate therapy in conjunction with treatment with Ivermectin Paste. Reinfestation, and measures for its prevention, should also be considered. Consult your veterinarian if the condition does not improve.

Distributed by:
FARNAM COMPANIES, INC.
 P.O. Box 34820
 Phoenix, AZ 85067-4820 U.S.A.
www.ivercare.com
 © 2002, 2005, 2006, 2010
 Farnam Companies, Inc.
 Product Code: 100504274
 10-0851 • 99.0000263.00

Farnam, the Farnam logo, IverCare, Sure-Grip and red to yellow color gradation are trademarks of Farnam Companies, Inc. Trademarks of Central Garden & Pet Company

CENTRAL
 Garden & Pet

IVERCARE®
 (ivermectin) Paste 1.87%

For Oral Use In Horses Only
IVERCARE®
 (ivermectin) Paste 1.87%



Anthelmintic and Boticide
Removes worms and bots with a single dose.
 Contents will treat up to 1500 lb body weight

Treats up to 1,500 lbs

PROUDLY MADE IN THE USA

Net Wt. 0.26 oz (7.30 g)
 ANADA 200-390, Approved by FDA

IVERCARE®
 (ivermectin) Paste 1.87%

PROUDLY MADE IN THE USA

IVERCARE®
 (ivermectin) Paste 1.87%



NO VARNISH

Lot No. / Exp. Date:



PROUDLY MADE IN THE USA