



## Compendium of Veterinary Products - US edition

MERCK ANIMAL HEALTH

Intervet Inc.

2 GIRALDA FARMS, MADISON, NJ, 07940

Customer Service: 800-521-5767

Order Desk: 800-648-2118

Technical Service (Companion Animal): 800-224-5318

Technical Service (Livestock): 800-211-3573

Fax: 973-937-5557

Website: [www.merck-animal-health-usa.com](http://www.merck-animal-health-usa.com)



Every effort has been made to ensure the accuracy of the information published. However, it remains the responsibility of the readers to familiarize themselves with the product information contained on the USA product label or package insert.

### **Nobivac® Canine 1-DAPPv**

Intervet/Merck Animal Health

### **CANINE DISTEMPER-ADENOVIRUS TYPE 2-PARAINFLUENZA-PARVOVIRUS VACCINE**

#### **Modified Live Virus**

#### **FOR ANIMAL USE ONLY**

**IN THE ABSENCE OF A VETERINARIAN-CLIENT-PATIENT RELATIONSHIP, FEDERAL REGULATIONS PROHIBIT THE RELABELING, REPACKAGING, RESALE OR REDISTRIBUTION OF THE INDIVIDUAL CONTENTS OF THIS PACKAGE.**

THIMEROSAL FREE

#### **PRODUCT DESCRIPTION:**

Nobivac® Canine 1-DAPPv vaccine is a modified live virus vaccine for the vaccination of healthy dogs as an aid in the prevention of disease caused by canine distemper virus, adenovirus type 1 (hepatitis) and adenovirus type 2 (respiratory disease), canine parainfluenza virus, and canine parvovirus. Data indicate the development of corneal opacity is not associated with the use of this product.

## **DOSAGE AND ADMINISTRATION:**

Transfer contents of the sterile diluent vial to the Nobivac Canine 1-DAPPv vial aseptically. Mix gently until dissolved. Use entire contents immediately after rehydration.

Inject one dose (1 mL) subcutaneously or intramuscularly. The initial dose may be given at 6 weeks of age or older. Repeat at 2 to 4 week intervals until the dog is 12 weeks of age. A minimum of two doses is required for primary immunization. Annual revaccination with one dose is recommended. Following the advice of the American Veterinary Medical Association on Vaccination Principles, an appropriate revaccination program for individual animals should be made based upon veterinarian-client-patient relationships.

## **PRECAUTIONS:**

- Store at 2° - 7°C.
- Do not freeze.
- Do not mix with other vaccines.
- Use new, non-chemically sterilized needles and syringes.
- Burn vaccine container and all unused contents.

## **WARNINGS:**

- Contains gentamicin as a preservative.
- Vaccinate only healthy, non-parasitized dogs.
- Do not vaccinate pregnant bitches.
- The use of a biological product may produce anaphylaxis and/or other inflammatory immune-mediated hypersensitivity reactions. **Antidote:** Epinephrine, corticosteroids, and antihistamines may all be indicated depending on the nature and severity of the reaction.
- The age at which maternal antibody for canine parvovirus no longer interferes with the development of active immunity varies according to the bitch's titer and quantity of colostral antibodies absorbed by the puppy.

Intervet Inc., Merck Animal Health, division of Intervet Inc., Omaha, NE 68103 USA

U.S. Veterinary License No. 165A

1 800 224-5318 (USA)

1 866 683-7838 (Canada)

For patent information:

<http://www.merck.com/product/patent/home.html>

Code

25x1 DOSE 065288 147913-02

**CPN:** 1047082.5

By using this site, you agree to the [Terms of Service](#) and [Privacy Policy](#).

[Contact Us](#) - [Logout](#)

Updated March 23, 2018

Copyright 2018 - Animalytix LLC