



Compendium of Veterinary Products - US edition

MERCK ANIMAL HEALTH

Intervet Inc.

2 GIRALDA FARMS, MADISON, NJ, 07940

Customer Service: 800-521-5767

Order Desk: 800-648-2118

Technical Service (Companion Animal): 800-224-5318

Technical Service (Livestock): 800-211-3573

Fax: 973-937-5557

Website: www.merck-animal-health-usa.com



Every effort has been made to ensure the accuracy of the information published. However, it remains the responsibility of the readers to familiarize themselves with the product information contained on the USA product label or package insert.

Nobivac® Puppy-DPv

Intervet/Merck Animal Health

CANINE DISTEMPER-PARVOVIRUS VACCINE

Modified Live Virus

FOR ANIMAL USE ONLY

IN THE ABSENCE OF A VETERINARIAN-CLIENT-PATIENT RELATIONSHIP, FEDERAL REGULATIONS PROHIBIT THE RELABELING, REPACKAGING, RESALE OR REDISTRIBUTION OF THE INDIVIDUAL CONTENTS OF THIS PACKAGE.

THIMEROSAL FREE

PRODUCT DESCRIPTION:

Nobivac® Puppy-DPv is a modified live virus vaccine containing attenuated strains of canine distemper virus (CDV) and parvovirus (CPV). This vaccine is for vaccination of healthy dogs as an aid in the prevention of disease caused by canine distemper virus and canine parvovirus.

DOSAGE AND ADMINISTRATION:

Aseptically reconstitute the desiccated vaccine with the sterile diluent provided and administer 1mL by the subcutaneous route or intramuscular route. Initial vaccination of healthy dogs may be given as early as 4 weeks of age, with booster injections administered every 3-4 weeks until the dog is 12 weeks of age. Dogs over 12 weeks of age should initially receive 2 doses 3-4 weeks apart. Annual revaccination with one dose is recommended.

Following the advice of the American Veterinary Medical Association on Vaccination Principles, an appropriate revaccination program for individual animals should be made based upon veterinarian-client-patient relationships.

PRECAUTIONS:

- Store at 2° - 7°C (35°-45°F).
- Do not freeze.
- Shake well before use.
- Use entire contents immediately after rehydration.
- Use new, non-chemically sterilized needles and syringes.
- Burn vaccine container and all unused contents.

WARNINGS:

- Contains gentamicin as a preservative.
- Vaccinate only healthy, non-parasitized dogs.
- Do not vaccinate pregnant bitches.
- The use of a biological product may produce anaphylaxis and/or other inflammatory immune-mediated hypersensitivity reactions. **Antidote:** Epinephrine, corticosteroids, and antihistamines may all be indicated depending on the nature and severity of the reaction.

Intervet Inc., Merck Animal Health, division of Intervet Inc., Omaha, NE 68103 USA

U.S. Veterinary License No. 165A

1 800 224-5318 (USA)

1 866 683-7838 (CAN)

For patent information:

<http://www.merck.com/product/patent/home.html>

Code

25x1 DOSE 053260 149225-03

CPN: 1047353.4

By using this site, you agree to the [Terms of Service](#) and [Privacy Policy](#).

[Contact Us](#) - [Logout](#)

Updated March 23, 2018

Copyright 2018 - Animalytix LLC