

MERCK ANIMAL HEALTH

Intervet Inc.

2 GIRALDA FARMS, MADISON, NJ, 07940

Customer Service: 800-521-5767

Order Desk: 800-648-2118

Technical Service (Companion Animal): 800-224-5318

Technical Service (Livestock): 800-211-3573

Fax: 973-937-5557

Website: www.merck-animal-health-usa.com



Every effort has been made to ensure the accuracy of the information published. However, it remains the responsibility of the readers to familiarize themselves with the product information contained on the USA product label or package insert.

SAFE-GUARD® DEWORMER FOR BEEF & DAIRY CATTLE AND GOATS SUSPENSION
10%

Intervet/Merck Animal Health

(fenbendazole)

(100 mg/mL)

RESIDUE WARNINGS: Cattle must not be slaughtered within 8 days following treatment. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal. Goats must not be slaughtered for food within 6 days following treatment. Because a withdrawal time in milk has not been established, do not use in lactating goats. For dairy cattle, there is no milk withdrawal period.

Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism.

Keep this and all medication out of the reach of children.

INDICATIONS:

Beef and Dairy Cattle - 2.3 mg/lb (5 mg/kg) body weight for the removal and control of: Lungworm (*Dictyocaulus viviparus*); Stomach worm (adults): brown stomach worm (*Ostertagia ostertagi*); Stomach worms (adults and fourth stage larvae): barberpole worm (*Haemonchus contortus* and *H. placei*) and small stomach worm (*Trichostrongylus axei*); Intestinal worms (adults and fourth stage larvae): hookworm (*Bunostomum phlebotomum*), thread-necked intestinal worm (*Nematodirus helvetianus*), small intestinal worm (*Cooperia*

punctata and *C. oncophora*), bankrupt worm (*Trichostrongylus colubriformis*), and nodular worm (*Oesophagostomum radiatum*).

Goats - 2.3 mg/lb (5 mg/kg) body weight for the removal and control of: Stomach worms (adults): *Haemonchus contortus* and *Teladorsagia circumcincta*.

DIRECTIONS:

Determine the proper dose according to estimated body weight. Administer orally. The recommended dose of 2.3 mg/lb (5 mg/kg) of body weight is achieved when 2.3 mL of the drug are given for each 100 lbs body weight.

EXAMPLES:

Cattle:

Body Weight Amount

100 lb	2.3 mL
200 lb	4.6 mL
300 lb	6.9 mL
400 lb	9.2 mL
500 lb	11.5 mL
1,000 lb	23.0 mL
1,500 lb	34.5 mL

Goats:

Body Weight Amount

25 lb	0.6 mL
50 lb	1.2 mL
75 lb	1.7 mL
100 lb	2.3 mL
125 lb	2.9 mL

Under conditions of continued exposure to parasites, retreatment may be needed after 4 to 6 weeks.

Manufactured by: DPT Laboratories, San Antonio, TX 78215

Distributed by: **Intervet Inc.**, Millsboro, DE 19966

Store at or below 25°C (77°F). Protect from freezing. Shake well before use. Restricted drug (California) - use only as directed.

For Use in Animals Only.

NADA # 128-620, Approved by FDA

1,000 mL (33.8 fl oz) 697510-E, 697515-E

1 Gallon (3,785 mL) 697935-D, 697938-E

CPN: 1047396.1