



## Compendium of Veterinary Products - US edition

Triangle® 10 HB (BOEHRINGER INGELHEIM VETMEDICA, INC.)  
BOEHRINGER INGELHEIM VETMEDICA, INC.  
3239 SATELLITE BLVD., DULUTH, GA, 30096  
Telephone: 800-325-9167  
Fax: 816-236-2717  
Website: [www.bevaccinesmart.com](http://www.bevaccinesmart.com)  
Website: [www.bi-vetmedica.com](http://www.bi-vetmedica.com)  
Website: [www.haveweseenyourcatlately.com](http://www.haveweseenyourcatlately.com)  
Website: [www.metacam.com](http://www.metacam.com)  
Website: [www.mycathasdiabetes.com](http://www.mycathasdiabetes.com)  
Website: [www.prrsresearch.com](http://www.prrsresearch.com)  
Website: [www.prozinc.us](http://www.prozinc.us)  
Website: [www.vetera-vaccines.com](http://www.vetera-vaccines.com)  
Website: [www.vetmedin-us.com](http://www.vetmedin-us.com)  
Website: [www.yourdogsheart.com](http://www.yourdogsheart.com)  
Email: [info@productionvalues.us](mailto:info@productionvalues.us)



Every effort has been made to ensure the accuracy of the information published. However, it remains the responsibility of the readers to familiarize themselves with the product information contained on the USA product label or package insert.

### **TRIANGLE® 10 HB**

Boehringer Ingelheim

#### **Bovine Rhinotracheitis-Virus Diarrhea-Parainfluenza 3-Respiratory Syncytial Virus Vaccine, Killed Virus, *Leptospira Canicola*-Grippotyphosa-Hardjo-Icterohaemorrhagiae-Pomona Bacterin**

*with Enhance™*

**Indications:** For vaccination of healthy cattle as an aid in the prevention of disease caused by bovine virus diarrhea (BVD) types 1 & 2, parainfluenza 3 (PI3), and bovine respiratory syncytial virus (BRSV), leptospirosis caused by *Leptospira pomona*, *L. hardjo*, *L. grippotyphosa*, *L. canicola*, and *L. icterohaemorrhagiae*, prevention of urinary shedding of *L. borgpetersenii* serovar Hardjo in heifers 10 months of age or older, and as an aid in the reduction of clinical disease due to infectious bovine rhinotracheitis (IBR). Vaccinated animals subsequently exposed to *L. borgpetersenii* serovar Hardjo (type hardjo-bovis) have been shown to clear renal infections within 8 weeks of exposure.

**Composition:** Contains IBR, BVD types 1 and 2, PI3, and BRSV killed viruses, *L. pomona*, *L. hardjo*, *L. grippotyphosa*, *L. canicola*, and *L. icterohaemorrhagiae*, with neomycin, polymyxin B, and thimerosal as preservatives.

**Dosage and administration:** Shake well. Inject one 5 mL dose intramuscularly or subcutaneously using aseptic technique. Repeat in 14 to 28 days. A 5 mL booster dose is recommended annually or prior to time of stress or exposure. Calves vaccinated under six months of age should be revaccinated at six months. May be administered to pregnant animals at any stage of gestation. Protect animals from exposure for at least 14 days after the last dose of vaccine.

**Precautions:** Store in the dark at 35 - 45°F (2 - 7°C). Avoid freezing. Use entire contents when first opened. Do not vaccinate within 21 days before slaughter. In case of anaphylactoid reaction, administer epinephrine.

**Boehringer Ingelheim Vetmedica, Inc., St. Joseph, MO 64506**

**US Vet. Lic. No. 124**

10 Doses/50 mL 161201-03

50 Doses/250 mL 161202-03

**CPN:** 1028274.1

By using this site, you agree to the [Terms of Service](#) and [Privacy Policy](#).

[Contact Us](#) - [Logout](#)

Updated March 26, 2018

Copyright 2018 - Animalytix LLC