

• **READ ENTIRE LABEL BEFORE USE**

NET CONTENTS: One Quart (32 Fl Oz)

FIRST AID

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact the National Pesticide Information Center at 1-800-858-7378 for emergency medical treatment information.

If in Eyes:

- Hold eye open and rinse slowly and gently with water for 15–20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

If on Skin or Clothing:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15–20 minutes.
- Call a poison control center or doctor for treatment advice.

If Swallowed:

- Immediately call a poison control center or doctor.
- Do not induce vomiting unless told to do so by the poison control center or doctor.
- Do not give **any** liquid to the person.
- Do not give anything by mouth to an unconscious person.

If Inhaled:

- Move person to fresh air.
- If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.
- Call a poison control center or doctor for further treatment advice.

NOTE TO PHYSICIAN:

Contains petroleum distillate — vomiting may cause aspiration pneumonia.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING:

Causes substantial but temporary eye irritation. Do not get in eyes. Avoid contact with skin or clothing. Harmful if swallowed. Avoid breathing vapor. Wash thoroughly after handling. For *space* spraying indoors wear a dust/mist filtering respirator (MSHA/NIOSH approval number prefix TC-21C) or a NIOSH approved respirator with any R, P or HE filter.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to fish. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

PHYSICAL OR CHEMICAL HAZARDS

Combustible! Do not use or store near heat or open flame. Flashpoint minimum of 105°F (41°C).

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Pesticide Storage: Store and transport in an upright position. Store in a cool dry area inaccessible to children or pets.

Pesticide Disposal: *If partly filled*—Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

Container Disposal: *If empty*—Non-refillable container. Do not reuse or refill this container. Place in trash or offer for recycling if available.

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or for climatic modification and being grown in interior landscapes, ornamental gardens or parks, or on golf courses or lawns and grounds.

For use in non-food areas indoors and outside premises of hotels, motels, homes, preschools, nurseries, day camps, greenhouses, lathhouses, seed houses, food handling establishments, packaging and processing plants, commercial, industrial, office, institutional and storage buildings, dairies, milk storage rooms, egg processing plants, supermarkets, restaurants, bottling plants, canneries, flour and feed mills, bakeries, and hospitals. In hospitals, patients should be removed prior to treatment. Also for use in campers, mobile homes, buses, automobiles, trucks, and boxcars. Also for use in dog kennels and horse stables. In horse stables, not for application if horses will be used for food.

Do not use in food/feed areas of food/feed handling establishments, restaurants or other areas where food/feed is commercially prepared or processed. Do not use in serving areas while food/feed is exposed or facility is in operation. Serving areas are areas where prepared foods/feed are served such as dining rooms but excluding areas where foods/feed may be prepared or held. In the home, cover all food handling surfaces and cover or remove all food and cooking utensils or wash thoroughly after treatment. Not for use in federally inspected meat and poultry plants. Nonfood/feed areas are areas such as garbage rooms, lavatories, floor drains (to sewers), entries, vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets, and storage (after canning or bottling). Food/feed areas of commercial food/feed handling establishments where the product cannot be used include areas for receiving, serving, storage (dry, cold, frozen, raw), packaging (canning, bottling, wrapping, boxing), preparing (cleaning, slicing, cooking, grinding), edible waste storage, enclosed processing systems (mills, dairies, edible oils, syrups).

Do not apply this product in hospital patient rooms while occupied or in any rooms while occupied by the elderly or infirm. Do not apply in classrooms when in use. Do not apply to institutions (including libraries, sports facilities, etc.) when occupants are present in the immediate area.

DO NOT USE THIS PRODUCT IN OR ON ELECTRICAL EQUIPMENT DUE TO THE POSSIBILITY OF SHOCK HAZARD.

For contact and residual sprays, mix with water in the following manner:

FINISHED SPRAY	INSECT CONTROL (Fluid Ounces)	WATER
0.25%	2 ² / ₃ (5- ¹ / ₃ Tbsps.)	1 gallon
0.25%	² / ₃ (4 tsps.)	1 quart
0.50%	5 ¹ / ₂ (10 ² / ₃ Tbsps.)	1 gallon
0.50%	1- ¹ / ₃ (8 tsps.)	1 quart

For heavy infestations, use the 0.50% dilution. For lighter infestations and for maintenance purposes, use the 0.25% dilution. One gallon of finished spray is sufficient to treat 1,580 linear feet with a 6 inch spray band.

INSTRUCTIONS FOR USE ON ANIMALS AND ANIMAL PREMISES

CATTLE BARNS, SWINE HOUSES, MILK ROOMS, KENNELS, POULTRY HOUSES, HORSE BARNS, DAIRIES, MILKING PARLORS, FEED LOTS, STABLES, PADDOCKS.

For initial clean up of severe insect infestation, dilute at a rate of 1 part concentrate in 25 parts water (5.5 ounces per gallon) (0.5%). For normal infestations dilute 1 part concentrate in 133 parts water (1.0 ounces per gallon) (2 tablespoons per gallon) (0.1%).

REFERENCE CHART

Pests	Dilutions	How to Apply
PESTS ON FARM PREMISES (Barns, Dairies, Loafing Sheds, Milking Parlors, Feedlots, Stables, Paddocks, and Livestock Housing)		
Horseflies, Stableflies, Lesser Houseflies, and other Manure Breeding Flies, Fleas, Lice, Mites, Ticks, including Deer Tick. Aids in control of Cockroaches, Mosquitoes, and Spiders.	Dilute 8 oz. in 8.25 gal. water. Dilute 1 pt. in 16.5 gal. water. Dilute 1 qt. in 33 gal. water.	Spray walls and surfaces to the point of runoff, about 1 gal. per 750-1000 sq. ft. Treat no more than once every two weeks. Do not contaminate animal's feed or water by spray. Do not use in milk rooms.
PESTS ON DOG PREMISES (Kennels, Dog Houses, Runs and Yards)		
Fleas, Flies, Lice and Mange Mites (adult and larvae)	Dilute 8 oz. in 8.25 gal. water. Dilute 1 pt. in 16.5 gal. water. Dilute 1 qt. in 33 gal. water.	Thoroughly wet pet's bedding or resting areas to runoff. Most efficient pest control is accomplished when heavy pressure and good contact is achieved. Timing and frequency of applications should be based on pest populations reaching nuisance levels, but accompanying manure removal and sanitation practices should precede sprays. Do not spray feed, food, or water. Repeat as needed but not more often than once every two weeks.
PESTS IN CATTLE BARN, HORSE BARN, SWINE HOUSES, KENNELS, MILKING PARLORS, MILK ROOMS, DAIRIES, POULTRY HOUSES, FEEDLOTS, STABLES, AND PADDOCKS.		
Flies, Mosquitoes and Gnats	Dilute 8 oz. in 8.25 gal. water. Dilute 1 pt. in 16.5 gal. water. Dilute 1 qt. in 33 gal. water.	<p>FOG OR FINE MIST: Apply, directing the spray toward the ceiling and upper corners until the area is filled with mist. Apply as a fog or mist, directing the nozzle for maximum coverage of area. Use about 2 oz. per 1000 cubic feet of space. For best results, close doors and windows before spraying and keep closed for ten to fifteen min. Vacate the treated area and ventilate before reoccupying. Repeat as needed but not more than once every two weeks.</p> <p>FOR USE THROUGH AUTOMATIC SPRAYING SYSTEMS: Do not program automatic spraying system to release pesticides where people, animals, food or feed are directly exposed. Do not install within three (3) feet of air ducts. Refer to system user manual for specific instructions. Dilute as directed. Dispense diluted concentrate through automatic spraying system. Set nozzles to deliver two (2) ounces per 1,000 cubic feet of space. Set timer to operate in accordance with equipment directions.</p>

— Continued —

REFERENCE CHART

Pests	Dilutions	How to Apply
PESTS ON LARGE ANIMALS (Dairy or Beef Cattle, Horses, Sheep, and Goats)		
Faceflies, Hornflies, Horseflies, Ear Ticks	Dilute 8 oz. in 6.5 gal. oil. Dilute 1 pt. in 13 gal. oil. Dilute 1 qt. in 26 gal. oil.	BACK RUBBER: Recharge backrubber as needed. Spray lactating dairy cows only after milking is completed.
Faceflies, Hornflies, Stableflies, Mosquitoes, Lice, Mites, Ticks, including Deer Tick	Dilute 8 oz. in 66.5 gal. water. Dilute 1 pt. in 133 gal. water. Dilute 1 qt. in 266 gal. water.	Spray to thoroughly cover entire animal. For Lice or Mites, a second treatment is recommended 14-21 days later. Spray lactating dairy cows only after milking is completed.
Hornflies, Faceflies, Stableflies, Ear Ticks	Dilute 2 oz. in 1.3 gal. water. Dilute 1 pt. in 3.3 gal. water	SPOT TREATMENT – LOW PRESSURE SPRAY Apply 1/2 oz. per ear or 2-4 oz. on face or 12-16 oz. along the backline.
PESTS ON SWINE AND POULTRY (Do not ship swine for slaughter within 5 days of last treatment.)		
Mange Mites	Dilute 8 oz. in 33 gal. water. Dilute 1 pt. in 66 gal. water. Dilute 1 qt. in 132 gal. water.	Spray, dip or sponge animals. Repeat after 14 days, spraying walls and floor space and bedding to kill late hatching, developing stages.
Blowflies, Flies, Mosquitoes, Hog Lice, Ticks, including Deer Ticks	Dilute 8 oz. in 33 gal. water. Dilute 1 pt. in 66 gal. water. Dilute 1 qt. in 132 gal. water.	Spray, dip or sponge to apply 1 pt. per pig, especially around ears.
Poultry Mites, Northern Fowl Mites, and Lice	Dilute 8 oz. in 16.5 gal. water. Dilute 1 pt. in 33 gal. water. Dilute 1 qt. in 66 gal. water.	Spray at the rate of 1/2 oz. per bird, or 1 gal. per 100 birds, with a fine mist. Spray roosts, walls and nests or cages.
PESTS ON DOGS (READ ENTIRE LABEL BEFORE EACH USE.) USE ONLY ON DOGS. Do not repeat treatment for seven (7) days. Do not use on puppies under twelve (12) weeks. Consult a veterinarian before using this product on debilitated, aged, pregnant, nursing or medicated animals.		
Fleas, Ticks, including Deer Tick, Lice and Mange	Dilute 8 oz. in 16.5 gal. water. Dilute 1 pt. in 33 gal. water. Dilute 1 qt. in 66 gal. water.	Thoroughly wet the animal by dipping, sponging or spraying. Allow animal to dry in a warm place without rinsing or toweling. Gives long lasting protection against reinfestation. Apply at intervals of 2 to 3 weeks. Sensitivities may occur after using ANY pesticide product for pets. If signs of sensitivity occur, bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately. In case of an emergency, call 1-800-548-2423 (Animal Poison Control Center).

INDOOR PEST CONTROL

For residual and contact control of cockroaches (waterbugs), crickets, silverfish (firebrats), clover mites, millipedes, centipedes, earwigs, sowbugs (pillbugs), spiders, brown dog ticks, lone star ticks, ants, fire ants, clothes moths, carpet beetles, fleas, boxelder bugs, scorpions, spider beetles, bedbugs, grasshoppers, house flies, mosquitoes, gnats, fruit flies, cluster flies, flying moths, confused flour beetles, saw-toothed grain beetles, shiny spider beetles, grain mites, red flour beetles, angoumois grain moths, grain weevils, Indian meal moths, Mediterranean flour moths, lesser grain borers, cigarette beetles, drug store beetles, chocolate moths. Spray thoroughly into hiding places such as cracks, crevices, moist areas, openings around pipes and sinks, under refrigerators, baseboards and storage areas. Spray directly on insects where possible.

— Continued on reverse —

For Bedbugs:

Apply as a spot treatment to cracks and crevices, around baseboards, floorboards, bedboards and walls. First, thoroughly clean and air mattresses and springs. Then spot treat only areas that may harbor bedbugs, concentrating on tufts, folds and seams. Allow all treated articles to dry thoroughly before use. Cover mattress with mattress cover or sheet before use. Do not apply to pillows or bed linens.

Grain Insects:

To control confused flour beetles, saw-toothed grain beetles, spider beetles, shiny spider beetles, grain mites, red flour beetles, angoumois grain moths, grain weevils, Indian meal moths, Mediterranean flour moths, lesser grain borers, cigarette beetles, drug store beetles, chocolate moths, carpet beetles — remove infested grain, cereal or flour and destroy. Avoid contamination of other food and foodstuffs in the areas by removing before spraying. Spray directly on exposed stages and into cracks and crevices where grain insects may be hiding. In homes, dispose of all infested products. The shelves and drawers should be cleaned and thoroughly sprayed so as to contact all pests, and covered with shelf paper after allowing shelves to dry. Any utensils that are contacted with spray should be washed before using.

Spiders:

Spray directly on these pests and to the webbing and all cracks and crevices which harbor spiders.

Ants:

To kill ants, spray trails, nests and points of entry within buildings and homes. Spray directly on ants where possible.

Wasps, Hornets, Yellow Jackets, Bees, Indoors:

Application should be made in the late evening when insects are at rest. Spray liberally into hiding and breeding places, especially under attic rafters, contacting as many insects as possible.

House Flies, Mosquitoes, Cluster Flies, Gnats, Fruit Flies, Flying Moths:

Close off all doors and windows. Point sprayer upward. Direct spray mist to all parts of the room, especially windows and other light sources which may attract these insects. Fill the room with mist, then leave the treated area. Keep the room closed for at least 15 minutes. Do not remain in treated room. Ventilate the room when treatment is completed.

Carpet Beetles and Clothes Moths:

To treat carpets — hold sprayer 14-18 inches from carpet. Apply spray evenly, covering the entire surface until slightly moist. Spray underside of carpet when possible. Avoid wetting or soaking. To treat Garments, Blankets, Carpets and Other Fabrics Before Storage — Dry clean or wash before treatment and after storage. Apply spray evenly, covering the material on both sides until slightly moist. For best results, place treated materials in plastic bags after drying, and store in a dark place. Also spray floor and baseboards, closets, cracks, crevices, infested areas of shelving and/or wherever these insects are seen or suspected.

NOTE:

When treating upholstery, carpeting, fabric, or finely finished surfaces, apply to an inconspicuous area first to observe any possible adverse reaction.

OUTDOOR PEST CONTROL

Pests on Outside Surfaces of Buildings:

Use this product to control ants, clover mites, crickets, earwigs, millipedes and sowbugs (pillbugs) on the outside surfaces of buildings. To help prevent infestation of buildings, treat the building foundation to a height of 2 to 3 feet where pests are and may find entrance. Also, apply as a residual spray to outside surfaces of buildings, including porches, screens, window frames, eaves, patios, garages, refuse dumps, and other areas where these pests congregate or have been seen.

House Flies, Mosquitoes, Cluster Flies, Gnats, Flying Moths, Outdoors:

For outdoor use only as an aid in reducing annoyance from these insects. Spray outside surfaces of screens, doors, window frames or wherever these insects may enter the room. Also treat surfaces around light fixtures on porches, in garages, and other places where these insects alight or congregate.

DIRECTIONS FOR USE IN OUTDOOR RESIDENTIAL MISTING SYSTEMS:

When using this product, installers and service technicians must comply with the license, certification, or registration requirements of the state(s), tribe(s) or local authority(ies) where they are installed. Do not apply this pesticide when people, pets and food are present. Do not use in an evaporative cooling system.

Apply as an area spray using an automatic spray system. Refer to the system user manual for specific instructions. Direct nozzles toward target area and away from swimming pools, water bodies, or eating and cooking areas. Set nozzles to deliver a fine mist or fog at a delivery rate not to exceed $1\frac{1}{4}$ fl. oz. of diluted product per minute, with each nozzle covering 100 sq. ft. (approximately 1,000 cu. ft.). This product may only be used in systems calibrated to apply no more than 0.69 fl. oz. of product (diluted 1 part concentrate with 21 parts of water) per 1,000 cu. ft. per day (equivalent to 0.25 grams of permethrin per 1,000 cu. ft. per day).

If system operates on an automatic timer, set the timing for application when people, pets and food are unlikely to be present. If system works when a person operates the remote activation device, then application of this pesticide when people, pets and food are present is prohibited.

If used in a system with a reservoir tank for the end use dilution, the system reservoir tank must be locked. Securely attach the end use pesticide label and a dilution statement to the system reservoir tank in a weather protected area or plastic sleeve. The dilution statement must be stated as follows: This container holds 1 part Permethrin Concentrate to 21 parts water.

If used in a direct injection system, the pesticide container must be locked. Securely attach the end use label to the pesticide container in a weather protected area or plastic sleeve.

Ants, Deer and Lone Star Ticks, Crickets, Fleas, and Earwigs, Outdoors:

For treatment of localized infestations of these insects in outdoor areas where there are weeds or bushy non-crop areas, spray infested areas thoroughly. For ants, thoroughly wet hills and runways.

Wasps, Hornets, Yellow Jackets, Bees, Outdoors:

Application should be made in late evening when insects are at rest. Aim spray at nest openings in ground, bushes, and in cracks and crevices which may harbor nests, saturating nest openings and contacting as many insects as possible.

To Control Fire Ants:

Dilute 2 tablespoons of this product in 1 gallon of water. Apply 1 gallon of solution as a gentle rain to each fire ant mound using a sprinkler can. Thoroughly wet the mound and surrounding area to a 4 foot diameter. For best results, apply in cool weather, 65 – 80°F, or in early morning or late evening hours. Treat new mounds as they appear. Do not disturb the mound prior to treatment. If possible, treat all mounds in the vicinity, and treat all colonies which may not as yet have constructed a mound. Larger quantities of solution can be prepared for multiple mound treatments. Equipment to deliver the diluted product as a gentle rain at the rate of 1 gallon per mound is still required. Do not use equipment that produces a pressurized spray as pressurized sprays may disturb fire ants and cause migration, reducing product effectiveness.

Note:

Seller warrants that this product complies with the specifications expressed in this label. To the extent consistent with applicable law, Seller makes no other warranties, and disclaims all other warranties, express or implied, including but not limited to warranties of merchantability and fitness for the intended purpose. To the extent consistent with applicable law, Seller's liability for default, breach or failure under this label shall be limited to the amount of the purchase price. To the extent consistent with applicable law, Seller shall have no liability for consequential damages.

For information on pesticide products (including health concerns, medical emergencies, or pesticide incidents), call the National Pesticide Information Center at 1-800-858-7378.



FARMGARD™

Permethrin Concentrate



TOTAL PEST CONTROL CONCENTRATE

For Use On Listed Livestock:

*Beef & Dairy Cattle, Horses, Sheep, Goats,
Swine, Poultry and their Premises*

*Use for Dogs & Kennels
Back Rubber*

- Controls Insect Pests for Up to 4 Weeks
- Controls Listed Flying & Crawling Insects

ACTIVE INGREDIENT

*Permethrin(CAS #52645-53-1) 13.30%w/w

OTHER INGREDIENTS** 86.70%

TOTAL...100.00%w/w

**KEEP OUT OF REACH
OF CHILDREN
WARNING**

See Back/Side/Other Panels for Additional
Precautionary Statements and First Aid.

NET CONTENTS:
One Quart (32 Fl Oz)

EPA Reg. No. 4-350-9444
EPA Est. No. 4-NY-1

*Cis/trans isomers ratio: min. 35% (±)cis and max. 65% (±) trans.

**Contains petroleum distillates.

Manufactured for Waterbury Companies, Inc., Waterbury, CT 06705
Customer Service: 800-845-3495 • Technical Information: 985-878-6751

