

# Equi-Jec® 6

## Influenza, Herpesvirus, Eastern & Western Encephalomyelitis, West Nile Virus Killed Vaccine, Tetanus Toxoid

### Description

Equi-Jec® 6 is an equine Influenza, Herpesvirus, Eastern, Western Encephalomyelitis, West Nile Virus, Tetanus vaccine.

### Features

- Protects against 6 diseases in 1 vaccine
- Safe for pregnant mares
- Safe for horses as young as 4 months of age
- Contains a contemporary West Nile Virus strain
- Duration of immunity 12 months for West Nile Virus

### Benefits

- Fewer needle sticks mean less pain for horses
- Single dose pre-filled syringe

### Advantages

Equi-Jec® 6 protects against Sleeping Sickness, West Nile Virus, Influenza, Herpesvirus (Rhino) and Tetanus in one vaccine.

### Dosage & Administration

Shake well before use. Using aseptic technique, administer to horses intramuscularly with a 1 mL dose. Administer a second 1 mL dose intramuscularly in 3-4 weeks using a different injection site. Revaccinate annually and prior to anticipated exposure with a single 1 mL intramuscular dose. Consultation with a veterinarian is recommended.

### Precautions

Store at 35-46° F (2-8° C). Protect from freezing. Do not vaccinate within 21 days before slaughter. Do not mix with other products.

Antidote: Epinephrine

### Highlights

- Protects horses against 6 diseases in 1 vaccine.
- Safe for pregnant mares
- Safe for horses as young as 4 months of age
- Contains a contemporary West Nile Virus strain
- Duration of immunity 12 months for West Nile Virus

### Uses

For vaccination of healthy horses 4 months of age or older as an aid in the prevention of disease caused by Eastern and Western Encephalomyelitis, and Tetanus; as an aid in the reduction of respiratory disease caused by Equine Influenza Virus (EIV) (product contains Kentucky lineage (KY/95), Florida sublineage clade 1 (OH/03), and Eurasian NewMarket/2/93 (NM 2/93) EIV strains), and equine Herpesvirus type 1 (EHV-1) and type 4 (EHV-4); as an aid in the reduction of severity of clinical disease, and as an aid in the prevention of viremia and mortality caused by West Nile Virus (WNV). Safe for use in pregnant mares.

### Warning

This product has been tested under laboratory conditions and shown to meet all Federal standards for safety and efficacy in normal, healthy, immunocompetent animals. This level of performance may be affected by conditions such as stress, weather, nutrition, disease, parasitism, other treatments, individual idiosyncrasies, or impaired immunological competency. These factors should be considered by the user when evaluating product performance or freedom from reactions.

### Product Information

Name	Code	Size
Equi-Jec® 6	125335	1 mL syringe 1 dose



1 Dose/1 mL

# Encephalomyelitis- Rhinopneumonitis-Influenza- West Nile Virus Vaccine Tetanus Toxoid

Eastern & Western, Killed Virus, Tetanus Toxoid

## Equi-Jec® 6


For Use In Animals Only

**Indications:** For vaccination of healthy horses 4 months of age or older as an aid in the prevention of disease caused by eastern and western encephalomyelitis and tetanus; as an aid in the reduction of respiratory disease caused by equine influenza virus (EIV) (product contains Kentucky Lineage (KY/95), Florida sublineage clade 1 (OH/03), and Eurasian Newmarket/2/93 (NM 2/93) EIV strains), and equine herpesvirus type 1 (EHV-1) and type 4 (EHV-4); as an aid in the reduction of severity of clinical disease, and an aid in the prevention of viremia and mortality caused by west nile virus (WNV). Safe for use in pregnant mares. The duration of immunity (DOI) is at least 6 months for OH/03 and at least 12 months for WNV.



Ser.:

Exp.:



Equi-Jec® 6

Peel Here  
136209-01

**Directions and dosage:** Shake well before use. Using aseptic technique, vaccinate horses intramuscularly with a 1 mL dose. Administer a second 1 mL dose intramuscularly in 3-4 weeks using a different injection site. The need for annual booster vaccinations has not been established for this product; consultation with a veterinarian is recommended.

**Precautions:** Store at 35–46°F (2–8°C). Do not freeze. Do not vaccinate within 21 days before slaughter. Do not mix with other products. In case of anaphylactoid reaction, administer epinephrine. In case of human exposure, contact a physician.

**Preservatives:** Gentamicin and Amphotericin B.

Equi-Jec® is a registered trademark of Boehringer Ingelheim Vetmedica GmbH. Used under license.

© 2017 Boehringer Ingelheim Vetmedica, Inc. All rights reserved.

US Patent Nos. 7,309,598; 8,697,089; 8,821,889; and 9,517,259.

US Patent Nos. 7,572,620; 8,535,685; 8,784,838; 9,180,181; and 9,492,530, used under license.


**Manufactured/Distributed By:**

**Boehringer Ingelheim  
Vetmedica, Inc.**

**St. Joseph, MO 64506**

**Phone: 866-638-2226**

**VLN/PCN 124/4855.24**



Vaccination Site

Equi-Jec is a registered trademark of Boehringer Ingelheim Vetmedica GmbH. Used under license.

© 2017 Boehringer Ingelheim Vetmedica, Inc. All rights reserved. EQU-0367-EQVAC1217

**Boehringer Ingelheim Vetmedica, Inc.**  
**2621 North Belt Highway • St. Joseph, MO 64506-2002**  
**Information 800-325-9167 • Orders 800-325-9167 • Fax 816-236-2717**  
**www.bi-vetmedica.com**