

MERCK ANIMAL HEALTH

Intervet Inc.

2 GIRALDA FARMS, MADISON, NJ, 07940

Customer Service: 800-521-5767
Order Desk: 800-648-2118
Technical Service (Companion Animal): 800-224-5318
Technical Service (Livestock): 800-211-3573
Fax: 973-937-5557
Website: www.merck-animal-health-usa.com



Every effort has been made to ensure the accuracy of the information published. However, it remains the responsibility of the readers to familiarize themselves with the product information contained on the USA product label or package insert.

Prestige® 3 + WNV

Intervet/Merck Animal Health

ENCEPHALOMYELITIS - WEST NILE VIRUS VACCINE, EASTERN & WESTERN, KILLED VIRUS, KILLED FLAVIVIRUS CHIMERA, TETANUS TOXOID

A combination of inactivated, concentrated, adjuvanted, tissue culture origin equine encephalomyelitis viruses (Eastern and Western), West Nile virus - flavivirus chimera and tetanus toxoid.

INDICATIONS: This product has been shown to be effective for the vaccination of healthy horses 6 months of age or older against Eastern and Western encephalomyelitis viruses, tetanus and West Nile Virus. For more information regarding safety and efficacy data, go to productdata.aphis.usda.gov.

This product has been shown to be effective against encephalitis and viremia caused by West Nile Virus.

DOSE: For primary vaccination aseptically administer 1 mL intramuscularly, repeat with a single dose in 3 - 4 weeks. Historically, annual revaccination with this product has been recommended. The need for annual booster vaccination has not been established for this product. For advice on revaccination in the face of stress or an exposure, consult your veterinarian.

CAUTIONS: Store at 2°-8°C (35°-46°F). Shake well before using. Do not mix with other products. This product has not been tested in pregnant animals. Foals nursing immune-dams should be vaccinated when maternal antibody levels will allow active immunization. Do not vaccinate within 21 days before slaughter. Inject deep into the muscle only. Local reactions may occur at the injection sites in vaccinates, and most of these reactions resolve in 3 to 7 days following vaccination. Systemic reactions may also occur in animals following vaccination. If allergic reaction occurs, treat with epinephrine. Care should be used to avoid injecting your own fingers, hands, or other body parts. Accidental injection can cause serious local reactions. In case of human exposure, contact a physician.

ANTIBIOTIC: Gentamicin.

PRESERVATIVE: Thimerosal.

FOR ANIMAL USE ONLY

Intervet Inc. d/b/a Merck Animal Health, Omaha, NE 68103, USA

VLN 165A/PCN 48W5.R1

1 800 521-5767 (USA)

1 866 683-7838 (CAN)

For patent information:

<http://www.merck.com/product/patent/home.html>

		Code	
10 Doses	10 x 1 mL Syringes with Separate Sterile Needles	197924	158834-01
10 Doses	10 mL	178734	187055-01

CPN: 1047541.0

By using this site, you agree to the [Terms of Service](#) and [Privacy Policy](#).
Copyright 2018 - Animalytix LLC